Cities xxx (xxxx) xxx–xxx

Contents lists available at ScienceDirect

Cities

journal homepage: www.elsevier.com/locate/cities

Implications of climate variability and change on urban and human health: A review

Israel Ropo Orimoloye*, Sonwabo Perez Mazinyo, Ahmed Mukalazi Kalumba, Olapeju Yewande Ekundayo, Werner Nel

Department of Geography and Environmental Science, University of Fort Hare, Private Bag X1314, Alice 5700, Eastern Cape Province, South Africa

ARTICLE INFO

Keywords:
Climate variability
Climate change
Implication
Human health
Extreme heat
Diseases

ABSTRACT

Urban regions have exceptional attributes that leave their dwellers and properties vulnerable to climate variability and change. Global temperatures continue to change, reaching new levels almost every year for the past two decades. This review examines the scientific evidence on the impact of climate change on urban and human health. It identifies research progress and gaps in how human society may respond to, adapt to, and prepare for the related changes. However, the causes are debated; climate variability and change are real. Climate variability and disaster risk are one of the threats to human health that adversely reinforce each other. Better knowledge of the linkage between climate change, variability and extreme weather-related illness is needed and can aid strategies to reduce the vulnerability. However, from this comprehensive review, it can be suggested that increased temperature and radiation are one of the major cause of some heat-related diseases such as skin cancer, heat stroke, heart disease and diarrhea which might strongly influenced by extreme climate events. On the other hand, since the extreme heat-related illnesses occurrence is increasing alarmingly, prevention and control have become a preference in public health programs and other disease control agencies. The study also suggests that public health should be everybody's business. Furthermore, public health education concepts can improve by a broader understanding of the subjective factors that underlie risk-taking and precautionary when exposed to extreme weather events.

1. Introduction

The global climate assessments revealed that Africa is susceptible to climate variability (Howard-Grenville, Buckle, Hoskins, & George, 2014; Onyango, Sahin, Awiti, Chu, & Mackey, 2016). Studies have shown that frequent storm occurrences, extreme heat, drought and other climate-related events are projected to increase as a result of both natural and human-made activities (Handmer et al., 2012; Kuenzer & Renaud, 2012; Nadin, Opitz-Stapleton, & Yinlong, 2015). The recognition of risk posed by climate on human health has created thriving concern in the linkage between climate adaptation and health threat (Benedikter, Läderach, Eitzinger, Cook, & Bruni, 2013; Chen, Doherty, Coffee, Wong, & Hellmann, 2016; Venton & La Trobe, 2008) by inspecting in depth this interlink between climate variability, risk reduction and human well-being. This review explores and investigates the synergies between heat-related events and human health. Also, to take a broader view of how the risk of extreme weather affects human health and their wellbeing. Therefore, to understand the significance of these risks over the urban areas, how they affect the environment and the crucial basis of urban functions that can be inimical to human health should be assessed.

Reed et al. (2013) revealed that the pronounced impact of climate-related events on human health may increase in the nearest future although the interdependence of climate variability and extreme weather remains ambivalent. Better knowledge of this interdependence is required and this can help planning to reduce its vulnerability (Dube, Moyo, Ncube, & Nyathi, 2016; Reckien et al., 2014; Voskamp & Van de Ven, 2015). Climate variability has been identified to pose numerous risks to human health across the globe (Austin et al., 2015) especially Africa (Sewankambo, Tumwine, & Besada, 2016) including South Africa (Ropo, Perez, Werner, & Enoch, 2017; Wright et al., 2017). The effect of climate on the environment has been studied (Aswani, Vaccaro, Abernethy, Albert, & de Pablo, 2015) and cannot be overemphasized as it cuts across every area of human life including health, natural resources and agriculture (Orimoloye & Adigun, 2017; Orimoloye, Mazinyo, Nel, & Kalumba, 2018; Ropo et al., 2017).

Climate variability contribute to both human-induced and natural disasters such as floods, toxic emissions, global warming, nuclear

* Corresponding author.
E-mail address: orimoloyeisrael@gmail.com (I.R. Orimoloye).

https://doi.org/10.1016/j.cities.2019.01.009
Received 13 August 2018; Received in revised form 20 December 2018; Accepted 5 January 2019
0264-2751/ © 2019 Elsevier Ltd. All rights reserved.

ARTICLE IN PRESS

explosions, fires, heat waves, drought, earthquake and rise in sea surface temperature as a result of thermal expansion with their related population shift, overcrowding and congestion which may support disease outbreak (Corcoran, 2016; Infield, Abunnasr, & Ryan, 2018). Many of the world's bothersome diseases that have lingered or persisted for years and are extensively influenced by climatic features include rainfall trend, extreme temperature (heat), humidity and wind patterns (El Morjani, Ebener, Boos, Ghaffar, & Musani, 2007; Ropo et al., 2017). More so, the incidence of extreme weather-related illness such as heart disease and skin cancers among others have continued to increase (Chang et al., 2016; Daniel, 2015; Xiang, Bi, Pisaniello, & Hansen, 2014) which might be as a result of increased climate variability globally. Therefore, the above-mentioned heat-related illnesses had been explored in this study globally and especially African countries including South Africa due to the recent changing climate potential impacts on human health (Ropo et al., 2017).

Studies have shown that change in climate parameters such as humidity, temperature, radiation and the wind correlates with these extreme weather-related illnesses (Glaser et al., 2016; Matysiak & Roess, 2017). Heat-related diseases such as cardiovascular disease, diarrhea, sunburn, heat stroke and skin cancer are among the dreaded diseases that trouble the world and as some of them kill faster than Tuberculosis and HIV/AIDS globally both in the developed and the developing nations including South Africa (CANSA REPORT, 2016). Preparedness and prevention have become vital themes in public health, emergency management and systems (Moore et al., 2007). In public health, risk reduction and management should be the focal point of the health experts and scientists. The idea and knowledge of climate variability are powerful scientific information that plays a significant role in human health management and risk reduction (Landauer, Juhola, & Klein, 2018; Rosenzweig et al., 2018). However, this idea of the linkage between extreme weather events and human is yet to be well explored or underutilized (Nehama et al., 2016; Patz & Frumkin, 2016). Therefore, this paper explored various studies on climate events-related impact on human and environmental health using information retrieved from Google Scholar and reports. The keywords for the search engine are climate variability, climate risk perspective and disaster in urban areas, spatial assessment of urban land surface heat and human health risk assessment, extreme heat and associated diseases.

1.1. The reality of climate variability

There are several adoptable pieces of literature that justify the fact that climate variability is really occurring and more evidence and research-based are being done to define this phenomenon (Abatzoglou, Rupp, & Mote, 2014; Kay et al., 2015; Orimoloye, Mazinyo, Nel, & Iortyom, 2018; Otto-Bliesner et al., 2016). Reports have shown that Southern Africa is experiencing climate variability which is evidenced by the occurrence of floods, extreme heat and droughts that are also predicted to be more visible or intensify in the nearest future (Alexander, Carzolio, Goodin, & Vance, 2013; Stringer et al., 2009; Ziervogel et al., 2014). Scientific research has also shown that the rising global air temperatures have a significant effect on the human health, rising sea levels and the environment (Turner et al., 2010; Venton & La Trobe, 2008; Watson, 2014).

Studies have equally indicated that there are scientific models that are more significant in analyzing climate scenarios as justification for climate variability (Berry & Richardson, 2016; Di Falco & Veronesi, 2014). According to Rosenzweig and Iglesias (1994) cited in Calzadilla et al. (2013), their models have been identified that climate variability is evident on agricultural productivity and verified how climate variability may alter the agricultural yields (Orimoloye & Adigun, 2017). The experimental findings on the effects of climate variability on crop yield summarized in Challinor et al. (2014) have, therefore, revealed a significant increase or rising global temperatures due to a warming climate.

ActionAid-Malawi (2006) states that many socio-economic problems which the poor societies in some part of the world for example, Malawi and other African countries are experiencing difficulties attributed to extreme weather events that closely associated with climate variability and its adverse impacts on human health. It is further highlighted that the climatic hazards are threatening the sustainable livelihoods of the majority family in the low-income nations (ActionAid-Malawi, 2006; Islam, 2013; Yawson et al., 2015). Studies have shown that a significant number of human are vulnerable to climate variability and their associated health threats (Deressa, Hassan, & Ringler, 2009; Islam, 2013; Nguyen, Bonetti, Rogers, & Woodroffe, 2016; Opiyo, 2014). Therefore, from the literature, it is evident that climate variability is real.

1.2. Global climate variability and its evidence in South Africa

The global warming prediction has revealed that the world will continue to experience increased warming and extreme climate events including heat waves, temperature and extreme heat are expected to annex in both intensity and frequency (Kendon et al., 2014; Moritz & Agudo, 2013). Findings have given its scientific evidence on various climatic scenarios that affect African nations and its population (Bryan, Deressa, Gbetibouo, & Ringler, 2009; Gbetibouo, 2009). Firstly, it is revealed that some countries such as South Africa experience a wide range of extreme climatic events that directly affect people's health and their well-being particularly those from rural areas because of their high dependence on climate-sensitive livelihood activities such as farming (Islam, 2013; Senapati & Gupta, 2017; Yawson et al., 2015). The most notable extreme climate events that affect human include extreme heat, droughts, floods, landslides, heat waves and the strong winds which have in many times contributed to poverty, famine, disease outbreak and loss of life among the populations. For example, the devastating floods of 1994–1995 in Western Cape (Lukamba, 2010), floods of 2000 in Limpopo (Reason & Keibel, 2004), extreme heat (Kruger & Sekele, 2013) and Floods of 2011 in KwaZulu-Natal (Dlamini et al., 2011). Table 1 present noticeable extreme weather event in South Africa;

Elasha, Elhassan, Ahmed, and Zakieldin (2005) observed that climate variability will adversely affect socio-economic sectors of the world in which both rural and urban livelihoods are built, particularly in Africa. These sectors include “human health, water resources, agriculture, forestry, fisheries and human settlements, ecological systems” (Sonwa, Somorin, Jum, Bele, & Nkem, 2012; Tirado, Cohen, Aberman, Meerman, & Thompson, 2010). Elasha et al. (2005) have also identified developing countries such as Malawi, South Africa in Southern Africa to

Table 1
Evidence of impacts of climate disasters in South Africa 1975–2006.

Year	Form of disasters	Place	Source
1981	Floods	Laingsburg, Western Cape	Lukamba, 2010
2010	Floods	Western Cape	Lukamba, 2010
1994–1995	Drought	Western Cape	Lukamba, 2010
2000	Floods	Limpopo	Reason & Keibel, 2004.
2013	Extreme heat (heatstroke)	North West	Kruger & Sekele, 2013
2011	Floods	KwaZulu-Natal province	Dlamini et al., 2011

be more vulnerable to extreme climate events because most of these countries have livelihood activities that are susceptible to climate events (Islam, 2013; Senapati & Gupta, 2017; Yawson et al., 2015). Studies have also revealed that most nations are affected by extreme climatic events such as extreme heat, droughts, heat waves, floods, strong winds and landslides every rainy season (Easterling et al., 2000; Kundzewicz, 2016).

Furthermore, it was also projected that the impacts of future climate extreme on health, economies and livelihood assets of poor societies in developing countries, predominantly among the poor urban, will be very devastating if proper measures for mitigation and adaptation are not put in place (Basher et al., 2015; Satterthwaite, 2007). In addition, climate variability is likely to impact negatively on health, environment, agriculture and development agenda of developing countries such as poverty reduction as the associated events have direct effects on poor people's livelihoods and the assets that determine their existence (Basher et al., 2015; Ibarrarán, Ruth, Ahmad, & London, 2009).

1.3. Climate risk perspective and disaster in urban areas

World cities, especially in the developing nations are faced with enormous problems such as a potential decline in the economic and human resources resulted from natural woes aggravated by the rate of unhindered urban growth and climate variability (González, Monsalve, Moris, & Herrera, 2018; Tan et al., 2010). There are several environmental dilemmas varieties of impact on cities (Neira & Prüss-Ustün, 2016) and the citizens' livelihoods due to climate variability (González et al., 2018). Inhabitants of the poor suburb areas and informal settlements are more vulnerable to hazards due to living in a high-risk environment such as slums, shanty, faulty shelters, blighted zone and as well as limited access to some basic services (Blaikie, Cannon, Davis, & Wisner, 2014; Mahmood et al., 2014). Consequently, there are tendencies of disease outbreaks in such areas.

Studies also revealed that, in the urban areas, the future heat extremes will be aggravated by the phenomenon called urban heat island and will in turn significantly influence urban health and the welfare of urban dwellers negatively (González et al., 2018; Tan et al., 2010). However, these can also trigger the existing illnesses that are related to the extreme heat and have effects on the ecosystem in the urban areas. Urban areas, in most cases, are more prone to disaster risk perhaps due to congestion, urban expansion, industrialization and inadequate or improper management (Begum, Sarkar, Jaafar, & Pereira, 2014; Schipper, 2009). Moreover, disaster risk is patterned over a period of time in the consequence of a critical interaction between development dynamics that bring about hazards and vulnerability. Unplanned urban settings are high-risk regions as rapid urbanization is accompanied by changes in land use, deforestation and reduction in vegetation cover and changes in climate variables, as the natural land surface are replaced by concrete, asphalt and structure which can influence or modify urban climate (Balica, Wright, & van der Meulen, 2012; Jha, Bloch, & Lamond, 2012).

In urban risk assessment, the hazard can be described as a harmful and dangerous phenomenon. Sometimes, human activities can create hazards, the substance that may cause a health-related problem, injury, loss of properties, decline in income, environmental degradation, economic disruption and as well as the loss of life (Hallegatte, 2014). The impacts of climate variability and hazards vary significantly with important variance in spatial pattern and location. For instance, slums that are located at the inner core of a city and overcrowded especially in developing nations, face many health threats and are vulnerable to hazards as a result of the dilapidated structures, congestion, difficulty in relocating and getting services (El-Fadel et al., 2014; Shaw, 2018). In the same way, suburb slums face other challenges for example, narrow road, drainage problem, unplanned layout, poor services and closeness to the environmental hazard.

Cities have exceptional attributes that leave their dwellers and valuable assets vulnerable to climate variability hazards. Many big cities, especially in low-income nations are found in the areas along the coast, slums and shanty, that are prone to hazards (Gasper, Blohm, & Ruth, 2011; Shamsuddoha, Ullah, & Shahjahan, 2014) these locations usually place human health and the economy at risk of extreme weather-related illnesses and disease outbreaks such as cholera, non-cholera diarrhea and cardiovascular disease. But this is not applicable only to city centers even the entire urban health can be affected by the threat caused by climate variability. The environmental aftermath of climate event is one of those that directly or indirectly affect the behavior and physical fitness of the dwellers including health impacts, water dearth and livelihood (Alexander et al., 2013; Braga, Zanobetti, & Schwartz, 2001). Climate extreme event is seen to be slow and advancing challenges combined with causes from multiple sources that are influenced by both anthropogenic and natural activities (Alexander et al., 2013; Anderegg, Prall, Harold, & Schneider, 2010).

Urban areas are the hotspots for identifying climate event risks. As a result of potential extreme weather exposure and climate-related illnesses, more than half of the city dwellers worldwide resided in the hazardous zones including congested zones, shanty, blighted areas and coastal areas (McGranahan, Balk, & Anderson, 2007). On the hand, urban regions also host the aforementioned crucial roles for wider socio-economy systems and often described by deep-rooted instability and vulnerability particular in big cities of the developing nations (Bull-Kamanga et al., 2003; Kraas, 2008). *Per contra*, climate change is not only peculiar to urban areas, but its impact also extends to the small cities and rural areas where human health, agriculture and economy are equally affected by the extreme weather due to climate variability.

Satterthwaite (2007) opined that inadequate attention has been given to the climate variability in the city areas and also to the extreme weather exposure of the city's population. In a contrary opinion, Mutunga and Hardee (2010) under the canopy of National Adaptation Programmes of Action (NAPA) reveals that almost all the focus of NAPA is on rural areas initiatives. But some of the developing countries are characterized by large numbers of people residing in rural areas. It is, however, noteworthy that these initiatives encompass strategies are few focused on climate and its effect on human well-being in the urban areas (Nkonya, Place, Kato, & Mwanjololo, 2015). Nevertheless, urban environment is home to the hotspot for a growing world's population. In this regards, full attention and concern of the urban environment are crucial in climate change implications on human health.

Moreover, urban areas comprehensively add to ecological degradation that ensnares significant risks of climate variability impact (Nkonya et al., 2015). The further amelioration of urban mitigation approach is a key obligation in managing the risks that posed by the extreme climate events and its adverse effect on the human health and environment. Additionally, more attention should be on the uniqueness or peculiarities of urban areas which frequently denote different scales, particularly in terms of built-up areas, population density and size, infrastructure, environmental degradation and rapid development compared to the rural areas (Carter et al., 2015; Tietenberg & Lewis, 2016).

The climate-related breakdown of the urban areas (big cities in particular) would imply challenges at broader scales beyond the city size (Kraas, 2008). But sometimes, the environmental problems or challenges of a particular area have nothing to do with the size of the city, for example, a city that is relatively small in size but located in the hazards prone. However, it is, therefore, essential to observe not only the bigger systems revolving around the processes or function depicted in the urban. But also, that the urban depend on the functions of environmental degradation surrounding it, as a result of climate events and its adverse indications on human health.

1.4. Geospatial appraisal of urban land surface heat and human health risk assessment

There is a developing acknowledgment in the fields of bio-meteorology and environmental health which posit that heat risk in urban

ARTICLE IN PRESS

Table 2
HI algorithms used in extreme weather events and its impact on human and environmental health.

No	Algorithm	Findings/gaps	References
1	$HI_c = T - 1.0799e^{0.03755T}_{c}\,(1 - e^{0.0801(D-14)}_{c})$	The model work perfectly at very low temperature but NWS model gives unrealistic value.	Schoen, 2005
2	$HI = -42.379 + 2.04901523T_f + 10.14333127H - 0.22475541T_f H - (6.83783 \times 10–3)\ T_f^2 - (5.481717 \times 10^{-2})\ H^2 + (1.22874 \times 10^{-3})\ T_f^2 H + (8.5282 \times 10^{-4})\ T_f H^2 - (1.99 \times 10^{-6})\ T_f^2 H^2.$	This is more suitable when temperatures are above 80 °F and relative humidity above 40%.	El Morjani et al., 2007
3	$HI_c = -2.719 + 0.994T_c + 0.016D_c^2$. Correction factor: $HIC = TC$ when $TC < 25$ °C	Applicable when temperatures are greater and equal to 25 °C.	Smoyer-Tomic & Rainham, 2001
4	$HI = 8.784695 + 1.61139411\,T + 2.338549RH - 0.14611605\,T * R\,H + 1.2308094 * 10^{-2}T^2 - 1.6424828 * 10^{-2}\,R\,H^2 + 2.211732 * 10^{-3}T^2\,R\,H + 7.2546\,10^{-4} * T * R\,H^2 - 3.582 * 10^{-6} * T^2$	This is valid only for air temperatures above 20 °C and its values are categorized due to possible heat perceived by human body.	Blazejczyk et al., 2012
5	$Hmidex(°C) = T_a + 0.555\,(e - 10)$ where $a = 6.11 * e^{(5417.7530*(\frac{1}{273.16} - \frac{1}{dew\ point + 273.16}))}$	Only endorsed for workplaces that have no "hot processes" as the main source of heat.	Jay and Kenny, 2010
6	$HI_c = T_C - 0.55 \times (1 - 0.001\ H)\ (T_C - 14.5)$	Recommended for both indoors and outdoors studies but may vary geographically and with locations.	Perry et al., 2011
7	$HI = K_1 + K_2\,T + K_3R + K_4TR + K_5\,T^2 + K_6R^2 + K_7T^2R + K_8TR^2 + K_9T^2R^2 + K_{10}\,T^3 + K_{11}R^3 + K_{12}T^3R + K_{13}TR^3 + K_{14}T^3R^2 + K_{15}T^2R^3 + K_{16}T^3R^3$	Recommended for both indoors and outdoors studies and this also vary with climatic parameters and locations.	Stull, 2000; Ropo et al., 2017

areas is a serious concern globally for example, USA (Basu & Samet, 2002; O'Neill & Ebi, 2009), South Africa (Orimoloye, Mazinyo, Nel, & Kalumba, 2018; Ropo et al., 2017; Wright et al., 2017), Australia (Vaneckova, Hart, Beggs, & De Dear, 2008) and Asia (Honda, 2007). Land use and land cover change (LULCC) is the investigation of land surface changes. Land use defines anthropogenic activities on land, while land cover explains the biophysical characteristics of the land surface. Land use change oft-times influence land cover features, thermal characteristics while changing land cover can likewise affect land use. Urbanization is essential evidence of LULCC which is synonymous with the modification of the surface energy budget resulting in rising temperature which can, in turn, have a significant impact on human health (Ige, Ajayi, Adeyeri, & Oyekan, 2017; Ogunjobi, Adamu, Akinsanola, & Orimoloye, 2018; Tursilowati & Djundjunan, 2007).

Land surface thermal signals such as land surface temperature which modifies the air temperature of the lower layers of the atmosphere are of prime significance to the urban context due to its important role in the energy balance of the surface. Land surface temperature not only helps to determine the internal climate among buildings but also influences energy exchanges that affect the comfort of city residents (Ogunjobi et al., 2018; Quan et al., 2014; Voogt & Oke, 2003). Alteration of the natural landscape in urban regions has been identified to make both local air and surface temperatures to rise a few degrees higher than that of surrounding rural areas (Streutker, 2003). Extreme heat affects health when there is thermal discomfort, this occurs when the body becomes too hot and there is no thermal equilibrium between the environment and the human body (Xu, Hu, Guan, & He, 2017).

Local micro-climatic parameters such as wind, air temperature, land surface temperature and radiation are often affected by natural objects and anthropogenic activities including built up, asphalted roads, pavement present in an environment. The disparity in the absorptive and reflective capabilities of objects to radiation associated with the heterogeneity of their physical characteristics often leads to the variation of these climatic elements which might affect the mental satisfaction of people and health in general (Lo & Quattrochi, 2003; Tursilowati & Djundjunan, 2007).

The relationship between local microclimate pattern and human thermal sensation has led to the development of several bio-meteorological indices which have been used to define human thermal comfort level (Ige et al., 2017; Ogunjobi et al., 2018). However, remote sensing and GIS techniques are considered to assess the impact of land use and

ARTICLE IN PRESS

land change on thermal characteristics and its consequences on human health because of its adequacy to obtain LULCC influenced surface thermal signal changes which often reshape people's health. For urban planning and environmental protection purposes monitoring human comfort shifts over various land use classifications is very important. Remote sensing information has been widely used in mapping the influence of human's activities on the environment taking remote sensing and geographic information system as tools for over the last two decades (Adefisan, Bayo, & Ropo, 2015; Adeola et al., 2017; Ogunjobi et al., 2018; Onamuti, Okogbue, & Orimoloye, 2017; Stephen, Iortyom, Ropo, & Daniel, 2017).

1.5. Extreme weather exposure and heat index (HI)

It is well known that discomfort that occurred in warm weather depends on the degree of temperature and humidity available in the air. The linkage between mortality and temperature has been examined by some researchers (Curriero et al., 2002; Gasparrini et al., 2015; Guo et al., 2014; McMichael et al., 2008). But more still need to be done regarding the geographical location and climatic features. The climate in recent years is characterized by the unusual increased in the intensity of extreme weather events and these have numerous effects on human health and the environment. The health implications of climate extreme have become a focal point and considered a crucial issue among climate researchers (Anderson, Bell, & Peng, 2013).

Several indices have been employed in studying the human health implications of climate variability and extreme weather. Some of the studies used HI (United State weather service (Hartz, Golden, Sister, Chuang, & Brazel, 2012; Metzger, Ito, & Matte, 2010)), humidex (Canadian meteorologist; Smoyer-Tomic & Rainham, 2001), wet-bulb globe temperature (Kjellstrom, Holmer, & Lemke, 2009; Moran & Epstein, 2006) and apparent temperature (Anderson & Bell, 2009; Chung et al., 2009) to assess HI indications of climate variability on human health and other researchers have used different algorithm to assess HI although with varying shortcomings (Table 2). Most studies in assessing the extreme heat implications on human thermal comfort and health have used ambient temperature (maximum, minimum and average temperature) to investigate heat-related illnesses (Hajat, O'Connor, & Kosatsky, 2010; Lin, de Dear, & Hwang, 2011; Wang, Hamann, Spittlehouse, & Murdock, 2012; Yu, Vaneckova, Mengersen, Pan, & Tong, 2010). Moreover, extreme weather implications cannot be fully assessed with the temperature only but other indicators such as radiation, wind and relative humidity should be considered to give a precise effect on human health (Lung, Lavalle, Hiederer, Dosio, & Bouwer, 2013). Among these indicators, the immediate surrounding has more effects on wind and radiation for example, wind speed is reduced by building effects or trees around and solar radiation is affected by localizing events include cloudiness and visibility (Parsons, 2014). If these factors (wind and radiation) are to be considered as variables and they are most suitable for a location as they can vary significantly in values over a short distance. The other two factors which are popularly used include relative humidity and air temperature because they are not spatially varied and these can be utilized to give a signal of the thermal comfort status over a large area (Commonwealth of Australia, 2005; Queiroz, Naas, & Sampaio, 2005; Ropo et al., 2017).

Heat index methods are statistically obtained equations that are analyzed (Table 2) to recreate the figures or values in Steadman (1979). These methods are substitutes to Steadman's algorithm for public and environmental health study and the latter algorithms considered the shortcomings of the former algorithm and modified it to be more accurate and efficient. Also, these algorithms can adequately calculate a longer sequence of heat stress index values based on the relative humidity and air temperature data. Moreover, these methods are popularly used to appraise HI thresholds for public health studies although there are variations in the use of the algorithm across the studies (Table 2). In reviewing the environmental studies literature this review pinpoints several heat stress indices algorithm but some are closely related and the present study suggests some of the algorithms and listed their findings and gaps. In public health and HI study, easier HI methods are less complex (for example, Halonen, Zanobetti, Sparrow, Vokonas, & Schwartz, 2011; Zanobetti & Schwartz, 2005; Vaneckova et al., 2011; Smoyer-Tomic & Rainham, 2001; Barnett, Tong, & Clements, 2010). More complex methods or formula that had been used in climate studies for example (Fischer & Schär, 2010; Oka, 2011).

1.6. The incidence of heart disease and climate variability

Excessive extreme heat exposure creates risks of some extreme heat-related problems including heat stroke, heart failure and inhibitions to sustain physical activities (Jendritzky & Tinz, 2009; Sahu, Sett, & Kjellstrom, 2013). The mortality due to extreme heat-related varies significantly by locations and climate zone. Studies have shown that people who reside in colder cities are affected by increased temperature while the hotter city dwellers are affected by colder temperatures (Curriero et al., 2002; Shanklin, 2004). A positive relationship between extreme heat and mortality (heart disease) has also been revealed, especially among the minors and elderly people (Basu & Samet, 2002; McGeehin & Mirabelli, 2001; Ropo et al., 2017). However, extreme weather can have detrimental effects on human health in both hotter and colder cities because thermal characteristics vary and sometimes maybe a personal experience-dependent and can be different from one person to another at the same time and space (Basu & Samet, 2002; Ryti, Guo, & Jaakkola, 2015).

More so, in some cases, extreme heat-related mortality occur in people with existing illnesses which include heart attack, heart failure and stroke. Urban dwellers are vulnerable to this risk than those who reside in rural areas (Basu & Samet, 2002; McGeehin & Mirabelli, 2001) because of their environments which have low ventilation and high thermal characteristics. Meteorologically this process is called urban heat island, a process where city or metropolitan region is significantly warmer than its surrounding suburban areas as a result of human activities. However, this can affect human health (Ketterer & Matzarakis, 2014; Vargo, Stone, Habeeb, Liu, & Russell, 2016). Behavioral and physiological adaptation as well as public health awareness can play a vital role in extreme heat morbidity and mortality reduction (Weisskopf et al., 2002). If the extreme heat is properly managed, the risk posed by the climate change will be minimal and extreme weather-related mortality can change over time by improving health care, public awareness, a study of this kind and personal understanding of climate change-related issues.

Schwartz, Samet, and Patz (2004) researched the impacts of temperature and humidity on human health and discovered that the hospital's admission (heart disease cases) increased after a few days of exposure to extreme weather. It has also been documented that hospital's admission for heart disease in the cities during hot and cold weather increases monotonically with increased temperature (both hot and cold weather) before admission (Mitchell & Chakraborty, 2015). However, this, therefore, connotes that the extreme weather contributes greatly to disease occurrence including cardiovascular disease regardless of age group. Seasonal extremes of weather are linked with increased mortality and are associated with weather-related illnesses while extreme heat and high temperatures have been linked with death among the minors and the olds (Lorenzini, Nali, & Pellegrini, 2014; Mitchell & Chakraborty, 2015; Suomi, 2014). In one hand, both hot and cold temperature-related mortality are dependents of latitude and locations (Braga et al., 2001; Davis, Hondula, & Patel, 2016; Kinney et al., 2015; Shi, Kloog, Zanobetti, Liu, & Schwartz, 2015). Therefore, extreme weather and health are significantly linked, however, most of the study on weather-related illnesses focused mostly on extreme temperatures while the impacts of humidity on human health and mortality rate have received little or no attention, the relative humidity plays a key role in determining human thermal stress index (Ropo et al., 2017).

1.7. Linkage between climate change and diarrhea incidence

Diarrhea is among the major causes of morbidity and mortality, particularly in developing nations. Hashizume et al. (2007) investigated the relationship between non-cholera diarrhea and climate events and the result revealed that the cases of hospital admission increase with increased temperature, especially in those respective at lower sanitation, social and economic status. Some studies suggest that about 4 billion episodes of diarrhea occurred in every year of which > 90% of this incidence takes place in the low-income nations (Aaronson, Marsh, Guha, Schuur, & Rouhani, 2015; Von-Doellinger, Campos, Mendes, & Schramm, 2014).

Investigations have shown seasonal coincidence of diarrhea incidence with the rainfall period, however, ambient temperature could also strengthen the diarrhea occurrence (Bush et al., 2014; Carlton, Woster, DeWitt, Goldstein, & Levy, 2015). In one hand, elucidating the significant role of extreme weather on the incidence and conveyance of diarrhea disease could help by bringing a broad insight into the mechanisms of the periodic of the disease. Multiple weather variables may contribute to the increased occurrence and outbreak of gastrointestinal illness. The infectious gastrointestinal is one of the most prevalent diseases globally and its principal symptom is diarrhea which caused about 3 million deaths annually among people of age 5 years and below especially in developing nations (Onozuka, Hashizume, & Hagihara, 2010). In developed nations, the related mortality may be low but the rate of morbidity remains high. The most incidences of diarrhea and related illnesses are brief and most times do not require medical attention but the socio-economic concerns are significant due to the high incidence of the disease. In recent years, with rapid concerns about climate variability globally few works have studied on the linkage between change in weather variables and variation in the diarrhea incidences and other related illnesses (Levy, Woster, Goldstein, & Carlton, 2016; Mellor et al., 2016; Phung et al., 2016).

Assessing the diarrhea occurrence is difficult by several significant risk factors and the issues of complication even without considering climate events impact (Mellor et al., 2016). These complexities include local disease etiology, inadequate hospital records, individual physiology, environmental factors, individual genetics, personal savior, engineered infrastructure, also as well as geographical, political, cultural and socio-economic elements (Rehfuess & Bartram, 2014). Moreover, environmental factors impacts such as temperature, humidity and wind trend might be altered by many factors such as available infrastructure, social condition and anthropogenic activities.

The changes in climate have been identified to pose several fundamental threats to human fitness (McMichael, 2014; Soret et al., 2014). The observation of climate trend at regional level (such as rising temperatures) and illnesses among children revealed positive relationship in Kenya since 1975. This was also projected that the drying and warming climate will continue to unveil as the population and anthropogenic activities continue to increased (Grace, Davenport, Funk, & Lerner, 2012).

Epidemiological analyses have helped in building an impression of the linkage between climate-related diarrhea occurrences. Nevertheless, given the inescapable threat of climate variability, there is a need for various strategies or approach in understanding the hazard occurrence of diarrhea and to plan viable ways to deal with enhancing and the perils that can be postured by environmental change and unusual heat.

This review identified several analytical studies that considered epidemiological and climatic variables that were linked together in order to appraise the linkage between extreme weather (extreme heat) and extreme heat-born disease outbreaks or the epidemic incidence. Ten of these studies were presented in the tabular form to know the types of weather variables they considered and also the temporal reference of the study (Table 3). The findings identified in the different studies but their results were varied, highlighted the linkage between extreme heat driven force and extreme weather-borne infections.

Table 3
Climate change and extreme heat-borne diseases, continent, country, period of study, Author, Data, and Data source.

S/no	Climate change borne disease	Continent	Country/region	Duration	Author	Data source (epidemic and climatic data)	Data
1	Diarrhea	Asia	Taiwan	1996–2007 (11 years)	Chou et al., 2010	National Health Insurance Research data base and Taiwan Central Weather Bureau	Temperature, relative humidity, and rainfall and diarrhea incidence
2	Diarrhea	Asia	China	2004–2011 (7 years)	Phung et al., 2015	Can-Tho Preventive Medicine Centre and Southern Regional Hydro-Meteorological Centre	Average temperature (°C), daily rainfall (mm) and average relative humidity (%) and epidemiological data.
3	Cardiovascular, diarrhea, and respiratory disease	South America, Asia, London, Sao Paulo, and Delhi		1991–1994 (4 years)	Hajat, Armstrong, Gouveia, & Wilkinson, 2005	Epidemic data (World Bank) Climatic data (unspecified)	Daily minimum &maximum temperature, relative humidity, Rainfall
4	Diarrhea	Africa	Botswana	1974–2003 (30 years)	Alexander et al., 2013	Climatic Data (Meteorology Department and Climate Research Unit)	Rainfall, temperature, and pressure.
5	Cardiovascular disease	North America	United States	19,986–1994 (8 years)	Schwartz et al., 2004	Climatic data (nearest airport) epidemiological data (Health Care Financing Admin.)	Temperature and relative humidity
6	Non-cholera diarrhea	Asia	Bangladesh	1996–2002 (7 years)	Hashizume et al., 2007	Climatic data (Bangladesh Meteorological Department) and epidemic data (International Centre for Diarrhea Disease Research, Bangladesh)	Rainfall and temperature
7	Diarrhea	Oceania (technically not a continent)	South Pacific	1993–1998 (5 years)	Singh et al., 2001	International Research Institute for climate and South Pacific Epidemiological service	Temperature, rainfall
8	General	Africa	South Africa	2015–2016 (6 months)	Wright et al., 2017	Field work	Temperature and humidity
9	Skin diseases	Global	Global	1990–2010 (20 years)	Hay et al., 2014	Systematic review and WHO	General
10	Skin cancer	London, Barcelona, Canary Islands	Europe	1979–1992 (13 years) & (2005–2014)	Grigalavicius, Moan, Dahlback, & Juzeniene, 2016	Nimbus 7 satellite and Institute of Health and Welfare, New Zealand Cancer Registry	UV Radiation, cloud cover, surface albedo, snow cover

Moreover, most of the studies revealed that there exists a positive relationship between the increased heat and epidemic incidence, also some revealed negative association that is there is an increase in diseases outbreak with decreased heat or temperature.

1.8. Skin diseases and climate variability

Cancer refers to the disease that affects the human body in which cells divide abnormally without restraint and infect other tissues in the human body. It is inimical impacts on health stretch from physical ineffectuality to demise (Andersen, Hercogová, Wollina, & Davis, 2012; Lavda, 2010). There are several forms of cancer and most times they are named after the name of the affected organ for example, skin, breast, lung and prostate cancers. Cancer is among the leading causes of mortality both in developed and under-developed nations after heart disease killing millions annually (Portier et al., 2017; Seyfried, 2015). There are potential skin cancers of both indirect and direct implication of climate variability. Climate variability may lead to a higher ambient temperature which can result in extreme weather event and change the subsequent human health exposures (Macdonald, Mackay, Li, & Hickie, 2003). Climate variability is also forecasted such as the increase in temperature and extreme weather events which will have impacts on human skin, extreme heat exposure can damage the skin (Andersen et al., 2012; Hunt & Watkiss, 2011). Moreover, more study is needed to determine the possibility of these issues of extreme heat-related disease, the location, geographical areas, the group of the population and people that are likely to be affected and its health implication or the outcome that could result. Albeit, the definite causes of cancer in human are yet to be understood absolutely for all classes of cancer, cancer growth factors or evolution include environmental contaminants, pathogens, age, genetic and sex (Kovarik & Kovarik, 2016). Highlighting the difficulties of understanding the inception or causes of skin disease and its related ailments, the connection among climate variability and skin malignancy is a blend of reality and speculation and studies are required to fill in the break in what we know to be achieved.

One of the likely direct effects of climate on skin disease can be as a result of much exposure to the extreme heat and radiation which might result in skin cancer. Most times, extreme heat effects on human skin are dependent on individual skin color. Skin color is one of the most apparent of human polymorphism (Avise & Ayala, 2010; Jablonski & Chaplin, 2010). One of the purposes of this study is to provide an encyclopedic review of epidemiology, etiology and climate change-related disease includes skin cancer. A clear understanding of the etiological factors is a crucial step in the avoidance of skin cancer (Schrempf et al., 2016).

Studies have shown that the aggregate of lifetime exposure to both artificial-natural heats is linked with persistent skin disorder such as skin cancer. Several studies that are epidemiologically oriented prove the elemental role of ultra-violet radiation in the provenance of skin cancer (Armstrong & Kricker, 2001; De Gruijl, van Kranen, & Mullenders, 2001; Haluza, Simic, Höltge, Cervinka, & Moshammer, 2014; Schrempf et al., 2016) in which most of the hazards result to melanoma. Study on the incidence of skin cancer in Florida (Rouhani et al., 2010), Chile (Rivas, Araya, Caba, Rojas, & Calaf, 2011) and Spain (Aceituno-Madera, Buendía-Eisman, Olmo, Jiménez-Moleón, & Serrano-Ortega, 2011) and related illness has been associated with a location, altitude and latitude. Moreover, it has been noted that the personal extreme heat exposure firmly depends on the behavioral elements or factors and also construed by the span of exposure, sun angle, geometry, clothing, surface, hair cover an hour of the day (Seckmeyer et al., 2013; Weihs et al., 2013).

1.9. Climate variability and human responses

Climate variability has been investigated to poses profound, direct or indirect and well-known threats to the human health and environment (Turner et al., 2010; Venton & La Trobe, 2008; Watson, 2014). A substantial portion of earth's species is projected to be vulnerable to decimation due to the increased temperature and other related extreme weather events due to changing climate and variability. There is rapid awareness of responses and the degree of risk, in both ardent and active that human will tackle as properties and lives are impaired by climate variability (Field, 2012). Climate variability can affect human health and the biological communities both indirect and direct impact. The direct human distress of climate variability includes changes in rainfall pattern, increased temperature, acidify and rising oceans, loss of ecosystems habitat (Ho, Knudby, Xu, Hodul, & Aminipouri, 2016; Parmesan & Yohe, 2003; Walther, 2010). However, the indirect aspect of climate may also affect the variety of life and ecosystem, sometimes more than the immediate consequence of climate variability.

The indirect effects arise as a result of attempts to both adapt and mitigate as to reduce or manage the climate variability adverse effect (Böckmann, 2015; Rounsevell, Dawson, & Harrison, 2010; Turner et al., 2010). There are many ways to this, but two are very crucial in addressing the undesirable concerns of climate variability on human fitness and well-being, these include adaptation and mitigation. Adapting to climate variability is crucial in order to avert the nonessential effects and to manage or improve the ecosystem resilience to simple stress (IPCC, 2014). The adaptation part was further divided by IPCC and recognized that there are two forms of adaptation which are spontaneous or autonomous adaptation and the second societal or planned adaptation. Spontaneous adaptation is when habitats and species include the various responses to the already observed adverse effect of climate. And the latter includes human risk reduction, policy and management process anticipated at expediting adaptation.

Several studies on mitigation and adaptation tend to research on their specific domain and few studies analyze the trivial consequences including adaptation options and the effects of migration assessment of impacts of climate variability (Birkmann, Garschagen, Kraas, & Quang, 2010; Füssel & Klein, 2006). Climate variability also has enormous responses or consequences on agriculture, human health, water, environment, where we live, what we eat and even the air we breathe. Moreover, climate variability can cause increase or decrease in agricultural output. They are some areas that experience an increase in agricultural potency because of climate variability. For instances, increased winter temperature could strengthen agricultural efficiency (Trnka et al., 2011) additionally summer temperature can diminish farming produce (Elliott et al., 2014).

Studies have opined that individual and societies can reduce their risk or threats posed by the climatic conditions by putting in place mitigation and adaptation strategies that best sustain their livelihoods (Kim et al., 2017; Waas & Hugé, 2012). The impacts of climate variability on livelihoods and human health are manifested by extreme events (extreme heat), floods, droughts, erratic rains and other environmental factors as established by Lema and Majule (2009). Africa has been categorized as one of the most vulnerable regions to climate variability because of its geographical location, incompetent leaders, widespread poverty compounded by these recurrent extreme weather events which limit the individual and communities' capacity to effectively cope and adapt to these challenges (Lema & Majule, 2009).

Studies have also revealed that the consequences of climate variability are potentially more substantial for the poor in low-income nations than for those living in more developed countries (Haines, Kovats, Campbell-Lendrum, & Corvalán, 2006; Seaman, Sawdon, Acidri, & Petty, 2014). Moreover, it is argued that vulnerability to the impacts of climate variability is a function of exposure to environmental factors or variables, sensitivity to those variables and the adaptive capacity of the affected individual or communities.

Krantz (2001) proposed and defined sustainable livelihood as follow:

> Right off the bat, “A livelihood involves the capacities, resources (stores, assets, claims and get to) and exercises required for a

method for a living. A livelihood is economical when it can adapt to and recoup from stress and stuns, keep up or upgrade its capacities and resources and give manageable livelihood opportunities to the present and the next generation; and which contributes net advantages to different livelihood at the neighborhood and global levels and in the short and long term".

From the above illustration, the ability of livelihood to be able to effectively adapt or cope with and recover from environmental stressors and shocks associated with climate and environmental change include extreme events, droughts, floods and extreme heat is central to the definition of sustainable livelihoods and environment (Carney, 2003).

2. Summary

In summary, the studies on climate-related events and heat-related diseases have been reviewed in this paper. Human health, diverse global socio-economic change, demographics, environment and climate variability are linked, such as the increased prevalence of heat-related disease, the emergence of infectious disease and the persistence of health disparity. Moreover, all these can be initiated and influenced by climate variability. Undertaking prevention at the local level to manage and reduce risks that may arise from these global climate influences is essential. Therefore, extreme weather and health are significantly linked, this review noted that most of the study on weather-related illnesses focused mostly on extreme temperatures but the impacts of humidity on human health and mortality rate have received little or no attention. Moreover, relative humidity plays a key role in determining human thermal stress index; this review opines that in the subsequent study of climate impacts on human health and their thermal comfort index more attention should be on humidity and other climatic variables.

There is an urgent demand to recognize the human health implications of climate change and variability. This will need the adaptation of present policies and approaches, a focus on an individual, public health education, institutions, government, and increase the awareness and communication across different sectors.

Acknowledgements

We would like to thank the University of Fort Hare, Alice South Africa for creating an enabling environment for research.

Competing interests

The authors declare that they have no competing interests.

Ethics, consent and permissions

No applicable.

Funding

Not applicable.

References

Aaronson, E. L., Marsh, R. H., Guha, M., Schuur, J. D., & Rouhani, S. A. (2015). Emergency department quality and safety indicators in resource-limited settings: An environmental survey. *International Journal of Emergency Medicine, 8*(1), 39.

Abatzoglou, J. T., Rupp, D. E., & Mote, P. W. (2014). Seasonal climate variability and change in the Pacific Northwest of the United States. *Journal of Climate, 27*(5), 2125–2142.

Aceituno-Madera, P., Buendía-Eisman, A., Olmo, F. J., Jiménez-Moleón, J. J., & Serrano-Ortega, S. (2011). Melanoma, altitude, and UV-B radiation. *Actas Dermo-Sifiliográficas, 102*(3), 199–205.

ActionAid-Malawi (2006). Climate change and smallholder farmers in Malawi. Understanding poor people's experiences in climate change adaptation. http://www.actionaid.org.uk/doc_lib/malawi_climate_change_report.pdf.

Adefisan, E. A., Bayo, A. S., & Ropo, O. I. (2015). Application of geo-spatial technology in identifying areas vulnerable to flooding in Ibadan metropolis. *Journal of Environment and Earth Science, 5*(14), 153–166.

Adeola, A. M., Olwoch, J. M., Botai, J. O., Rautenbach, C. D., Kalumba, A. M., Tsela, P. L., & Nsubuga, F. W. N. (2017). Landsat satellite derived environmental metric for mapping mosquitoes breeding habitats in the Nkomazi municipality, Mpumalanga Province, South Africa. *South African Geographical Journal, 99*(1), 14–28.

Alexander, K. A., Carzolio, M., Goodin, D., & Vance, E. (2013). Climate change is likely to worsen the public health threat of diarrheal disease in Botswana. *International Journal of Environmental Research and Public Health, 10*(4), 1202–1230.

Anderegg, W. R., Prall, J. W., Harold, J., & Schneider, S. H. (2010). Expert credibility in climate change. *Proceedings of the National Academy of Sciences, 107*(27), 12107–12109.

Andersen, L. K., Hercogová, J., Wollina, U., & Davis, M. D. (2012). Climate change and skin disease: A review of the English-language literature. *International Journal of Dermatology, 51*(6), 656–661.

Anderson, B. G., & Bell, M. L. (2009). Weather-related mortality: How heat, cold, and heat waves affect mortality in the United States. *Epidemiology (Cambridge, Mass.), 20*(2), 205.

Anderson, G. B., Bell, M. L., & Peng, R. D. (2013). Methods to calculate the heat index as an exposure metric in environmental health research. *Environmental Health Perspectives, 121*(10), 1111.

Armstrong, B. K., & Kricker, A. (2001). The epidemiology of UV induced skin cancer. *Journal of Photochemistry and Photobiology B: Biology, 63*(1), 8–18.

Aswani, S., Vaccaro, I., Abernethy, K., Albert, S., & de Pablo, J. F. L. (2015). Can perceptions of environmental and climate change in island communities assist in adaptation planning locally? *Environmental Management, 56*(6), 1487–1501.

Austin, S. E., Ford, J. D., Berrang-Ford, L., Araos, M., Parker, S., & Fleury, M. D. (2015). Public health adaptation to climate change in Canadian jurisdictions. *International Journal of Environmental Research and Public Health, 12*(1), 623–651.

Avise, J. C., & Ayala, F. J. (2010). *Human skin pigmentation as an adaptation to UV radiation*.

Balica, S. F., Wright, N. G., & van der Meulen, F. (2012). A flood vulnerability index for coastal cities and its use in assessing climate change impacts. *Natural Hazards, 64*(1), 73–105.

Barnett, A. G., Tong, S., & Clements, A. C. A. (2010). What measure of temperature is the best predictor of mortality? *Environmental Research, 110*(6), 604–611.

Basher, R., Hayward, B., Lavell, A., Martinelli, A., Perez, O., Pulwarty, R., & Cutter, S. (2015). *Disaster risks research and assessment to promote risk reduction and management*.

Basu, R., & Samet, J. M. (2002). Relation between elevated ambient temperature and mortality: A review of the epidemiologic evidence. *Epidemiologic Reviews, 24*(2), 190–202.

Begum, R. A., Sarkar, M. S. K., Jaafar, A. H., & Pereira, J. J. (2014). Toward conceptual frameworks for linking disaster risk reduction and climate change adaptation. *International Journal of Disaster Risk Reduction, 10*, 362–373.

Benedikter, A., Läderach, P., Eitzinger, A., Cook, S., & Bruni, M. (2013). Addressing adaptation to support disaster risk reduction: A framework for supply chain inclusive adaptation to climate change. *Climate change and disaster risk management* (pp. 513–533). Springer Berlin Heidelberg.

Berry, P., & Richardson, G. R. (2016). Approaches for building community resilience to extreme heat. *Extreme weather, health, and communities* (pp. 351–388). Springer International Publishing.

Birkmann, J., Garschagen, M., Kraas, F., & Quang, N. (2010). Adaptive urban governance: New challenges for the second generation of urban adaptation strategies to climate change. *Sustainability Science, 5*(2), 185–206.

Blaikie, P., Cannon, T., Davis, I., & Wisner, B. (2014). *At risk: natural hazards, people's vulnerability and disasters*. Routledge.

Blazejczyk, K., Epstein, Y., Jendritzky, G., Staiger, H., & Tinz, B. (2012). Comparison of UTCI to selected thermal indices. *International journal of biometeorology, 56*(3), 515–535.

Böckmann, M. (2015). *Exploring the health context: A multimethod approach to climate change adaptation evaluation* (Doctoral dissertation)Staats-und Universitätsbibliothek Bremen.

Braga, A. L. F., Zanobetti, A., & Schwartz, J. (2001). The time course of weather-related deaths. *Epidemiology, 12*(6), 662–667.

Bryan, E., Deressa, T. T., Gbetibouo, G. A., & Ringler, C. (2009). Adaptation to climate change in Ethiopia and South Africa: Options and constraints. *Environmental Science & Policy, 12*(4), 413–426.

Bull-Kamanga, L., Diagne, K., Lavell, A., Leon, E., Lerise, F., MacGregor, H., & Satterthwaite, D. (2003). From everyday hazards to disasters: The accumulation of risk in urban areas. *Environment and Urbanization, 15*(1), 193–204.

Bush, K. F., O'Neill, M. S., Li, S., Mukherjee, B., Hu, H., Ghosh, S., & Balakrishnan, K. (2014). Associations between extreme precipitation and gastrointestinal-related hospital admissions in Chennai, India. *Environmental Health Perspectives, 122*(3), 249.

Calzadilla, A., Rehdanz, K., Betts, R., Falloon, P., Wiltshire, A., & Tol, R. S. (2013). Climate change impacts on global agriculture. *Climatic Change, 120*(1–2), 357–374.

CANSA (2016). *The Cancer Association of South Africa report*.

Carlton, E. J., Woster, A. P., DeWitt, P., Goldstein, R. S., & Levy, K. (2015). A systematic review and meta-analysis of ambient temperature and diarrhoeal diseases. *International Journal of Epidemiology, 45*(1), 117–130.

Carney, D. (2003). *Sustainable livelihoods approaches: Progress and possibilities for change*. London: Department for International Development.

Carter, J. G., Cavan, G., Connelly, A., Guy, S., Handley, J., & Kazmierczak, A. (2015). Climate change and the city: Building capacity for urban adaptation. *Progress in Planning, 95*, 1–66.

Challinor, A. J., Watson, J., Lobell, D. B., Howden, S. M., Smith, D. R., & Chhetri, N.

(2014). A meta-analysis of crop yield under climate change and adaptation. *Nature Climate Change, 4*(4), 287–291.
Chang, H., Xiang, C., Duan, C., Wan, Z., Liu, Y., Zheng, Y., & Shu, S. (2016). Study on the thermal performance and wind environment in a residential community. *International Journal of Hydrogen Energy, 41*(35), 15868–15878.
Chen, C., Doherty, M., Coffee, J., Wong, T., & Hellmann, J. (2016). Measuring the adaptation gap: A framework for evaluating climate hazards and opportunities in urban areas. *Environmental Science & Policy, 66*, 403–419.
Chou, W. C., Wu, J. L., Wang, Y. C., Huang, H., Sung, F. C., & Chuang, C. Y. (2010). Modeling the impact of climate variability on diarrhea-associated diseases in Taiwan (1996–2007). *Science of the Total Environment, 409*(1), 43–51.
Chung, J. Y., Honda, Y., Hong, Y. C., Pan, X. C., Guo, Y. L., & Kim, H. (2009). Ambient temperature and mortality: An international study in four capital cities of East Asia. *Science of the Total Environment, 408*(2), 390–396.
Commonwealth of Australia (2005). Bureau of Meteorology (ABN 92 637 533 532). http://www.bom.gov.au/info/thermal_stress/.
Corcoran, J. (2016). *Implications of climate change for the livelihoods of urban dwellers in Kiribati* (Doctoral dissertation)University of Waikato.
Curriero, F. C., Heiner, K. S., Samet, J. M., Zeger, S. L., Strug, L., & Patz, J. A. (2002). Temperature and mortality in 11 cities of the eastern United States. *American Journal of Epidemiology, 155*(1), 80–87.
Daniel, O. A. (2015). Urban extreme weather: A challenge for a healthy living environment in Akure, Ondo State, Nigeria. *Climate, 3*(4), 775–791.
Davis, R. E., Hondula, D. M., & Patel, A. P. (2016). Temperature observation time and type influence estimates of heat-related mortality in seven US cities. *Environmental Health Perspectives, 124*(6), 795.
De Gruijl, F. R., van Kranen, H. J., & Mullenders, L. H. (2001). UV-induced DNA damage, repair, mutations and oncogenic pathways in skin cancer. *Journal of Photochemistry and Photobiology B: Biology, 63*(1), 19–27.
Deressa, T. T., Hassan, R. M., & Ringler, C. (2009). *Assessing household vulnerability to climate change. Vol. 935*. Intl Food Policy Res Inst.
Di Falco, S., & Veronesi, M. (2014). Managing environmental risk in presence of climate change: The role of adaptation in the Nile Basin of Ethiopia. *Environmental and Resource Economics, 57*(4), 553–577.
Dlamini, P., Orchard, C., Jewitt, G., Lorentz, S., Titshall, L., & Chaplot, V. (2011). Controlling factors of sheet erosion under degraded grasslands in the sloping lands of KwaZulu-Natal, South Africa. *Agricultural Water Management, 98*(11), 1711–1718.
Dube, T., Moyo, P., Ncube, M., & Nyathi, D. (2016). *The impact of climate change on agro-ecological based livelihoods in Africa: A review*.
Easterling, D. R., Evans, J. L., Groisman, P. Y., Karl, T. R., Kunkel, K. E., & Ambenje, P. (2000). Observed variability and trends in extreme climate events: A brief review. *Bulletin of the American Meteorological Society, 81*(3), 417–425.
El Morjani, Z. E. A., Ebener, S., Boos, J., Ghaffar, E. A., & Musani, A. (2007). Modelling the spatial distribution of five natural hazards in the context of the WHO/EMRO Atlas of Disaster Risk as a step towards the reduction of the health impact related to disasters. *International Journal of Health Geographics, 6*(1), 8.
Elasha, B. O., Elhassan, N. G., Ahmed, H., & Zakieldin, S. (2005). Sustainable livelihood approach for assessing community resilience to climate change: Case studies from Sudan. *Assessments of impacts and adaptations to climate change (AIACC) working paper* (pp. 17). .
El-Fadel, M., Maroun, R., Quba'a, R., Mawla, D., Sayess, R., Massoud, M. A., & Jamali, I. (2014). Determinants of diarrhea prevalence in urban slums: A comparative assessment towards enhanced environmental management. *Environmental Monitoring and Assessment, 186*(2), 665–677.
Elliott, J., Deryng, D., Müller, C., Frieler, K., Konzmann, M., Gerten, D., & Eisner, S. (2014). Constraints and potentials of future irrigation water availability on agricultural production under climate change. *Proceedings of the National Academy of Sciences, 111*(9), 3239–3244.
Field, C. B. (2012). *Managing the risks of extreme events and disasters to advance climate change adaptation: Special report of the intergovernmental panel on climate change*. Cambridge University Press.
Füssel, H. M., & Klein, R. J. (2006). Climate change vulnerability assessments: An evolution of conceptual thinking. *Climatic Change, 75*(3), 301–329.
Gasparrini, A., Guo, Y., Hashizume, M., Lavigne, E., Zanobetti, A., Schwartz, J., & Leone, M. (2015). Mortality risk attributable to high and low ambient temperature: A multicountry observational study. *The Lancet, 386*(9991), 369–375.
Gasper, R., Blohm, A., & Ruth, M. (2011). Social and economic impacts of climate change on the urban environment. *Current Opinion in Environmental Sustainability, 3*(3), 150–157.
Gbetibouo, G. A. (2009). *Understanding farmers' perceptions and adaptations to climate change and variability: The case of the Limpopo Basin, South Africa. Vol. 849*. Intl Food Policy Res Inst.
Glaser, J., Lemery, J., Rajagopalan, B., Diaz, H. F., García-Trabanino, R., Taduri, G., & Jha, V. (2016). Climate change and the emergent epidemic of CKD from heat stress in rural communities: The case for heat stress nephropathy. *Clinical Journal of the American Society of Nephrology, 11*(8), 1472–1483.
González, D. P., Monsalve, M., Moris, R., & Herrera, C. (2018). Risk and resilience monitor: Development of multiscale and multilevel indicators for disaster risk management for the communes and urban areas of Chile. *Applied Geography, 94*, 262–271.
Grace, K., Davenport, F., Funk, C., & Lerner, A. M. (2012). Child malnutrition and climate in Sub-Saharan Africa: An analysis of recent trends in Kenya. *Applied Geography, 35*(1), 405–413.
Grigalavicius, M., Moan, J., Dahlback, A., & Juzeniene, A. (2016). Daily, seasonal, and latitudinal variations in solar ultraviolet A and B radiation in relation to vitamin D production and risk for skin cancer. *International Journal of Dermatology, 55*(1).
Guo, Y., Gasparrini, A., Armstrong, B., Li, S., Tawatsupa, B., Tobias, A., & Tong, S. (2014). Global variation in the effects of ambient temperature on mortality: A systematic evaluation. *Epidemiology (Cambridge, Mass.), 25*(6), 781.
Haines, A., Kovats, R. S., Campbell-Lendrum, D., & Corvalán, C. (2006). Climate change and human health: Impacts, vulnerability and public health. *Public Health, 120*(7), 585–596.
Hajat, S., Armstrong, B. G., Gouveia, N., & Wilkinson, P. (2005). Mortality displacement of heat- related deaths: A comparison of Delhi, Sao Paulo, and London. *Epidemiology, 16*(5), 613–620.
Hajat, S., O'Connor, M., & Kosatsky, T. (2010). Health effects of hot weather: From awareness of risk factors to effective health protection. *The Lancet, 375*(9717), 856–863.
Hallegatte, S. (2014). Trends in hazards and the role of climate change. *Natural disasters and climate change* (pp. 77–97). Springer International Publishing.
Halonen, J. I., Zanobetti, A., Sparrow, D., Vokonas, P. S., & Schwartz, J. (2011). Outdoor temperature is associated with serum HDL and LDL. *Environmental Research, 111*(2), 281–287.
Haluza, D., Simic, S., Höltge, J., Cervinka, R., & Moshammer, H. (2014). Connectedness to nature and public (skin) health perspectives: Results of a representative, population-based survey among Austrian residents. *International Journal of Environmental Research and Public Health, 11*(1), 1176–1191.
Handmer, J., Honda, Y., Kundzewicz, Z. W., Arnell, N., Benito, G., Hatfield, J., & Takahashi, K. (2012). Changes in impacts of climate extremes: Human systems and ecosystems. *Managing the risks of extreme events and disasters to advance climate change adaptation special report of the intergovernmental panel on climate change* (pp. 231–290). Intergovernmental Panel on Climate Change.
Hartz, D. A., Golden, J. S., Sister, C., Chuang, W. C., & Brazel, A. J. (2012). Climate and heat- related emergencies in Chicago, Illinois (2003–2006). *International Journal of Biometeorology, 56*(1), 71–83.
Hashizume, M., Armstrong, B., Hajat, S., Wagatsuma, Y., Faruque, A. S., Hayashi, T., & Sack, D. A. (2007). Association between climate variability and hospital visits for non-cholera diarrhoea in Bangladesh: Effects and vulnerable groups. *International Journal of Epidemiology, 36*(5), 1030–1037.
Hay, R. J., Johns, N. E., Williams, H. C., Bolliger, I. W., Dellavalle, R. P., Margolis, D. J., & Michaud, C. (2014). The global burden of skin disease in 2010: An analysis of the prevalence and impact of skin conditions. *Journal of Investigative Dermatology, 134*(6), 1527–1534.
Ho, H. C., Knudby, A., Xu, Y., Hodul, M., & Aminipouri, M. (2016). A comparison of urban heat islands mapped using skin temperature, air temperature, and apparent temperature (Humidex), for the greater Vancouver area. *Science of the Total Environment, 544*, 929–938.
Honda, Y. (2007). Impact of climate change on human health in Asia and Japan. *Global Environmental Research, 11*(1), 33.
Howard-Grenville, J., Buckle, S. J., Hoskins, B. J., & George, G. (2014). Climate change and management. *Academy of Management Journal, 57*(3), 615–623.
Hunt, A., & Watkiss, P. (2011). Climate change impacts and adaptation in cities: A review of the literature. *Climatic Change, 104*(1), 13–49.
Ibarrarán, M. E., Ruth, M., Ahmad, S., & London, M. (2009). Climate change and natural disasters: Acroeconomic performance and distributional impacts. *Environment, Development and Sustainability, 11*(3), 549–569.
Ige, S. O., Ajayi, V. O., Adeyeri, O. E., & Oyekan, K. S. A. (2017). Assessing remotely sensed temperature humidity index as human comfort indicator relative to landuse landcover change in Abuja, Nigeria. *Spatial Information Research, 25*(4), 523–533.
Infield, E. M. H., Abunnasr, Y., & Ryan, R. L. (2018). *Planning for climate change: A reader in green infrastructure and sustainable design for resilient cities*. Routledge.
Intergovernmental Panel on Climate Change (2014). *Climate change 2014–impacts, adaptation and vulnerability: Regional aspects*. Cambridge University Press.
Islam, M. M. (2013). *Vulnerability and adaptation of fishing communities to the impacts of climate variability and change: Insights from coastal Bangladesh*. University of Leeds.
Jablonski, N. G., & Chaplin, G. (2010). Human skin pigmentation as an adaptation to UV radiation. *Proceedings of the National Academy of Sciences, 107*(Supplement 2), 8962–8968.
Jay, O., & Kenny, G. P. (2010). Heat exposure in the Canadian workplace. *American journal of industrial medicine, 53*(8), 842–853.
Jendritzky, G., & Tinz, B. (2009). The thermal environment of the human being on the global scale. *Global Health Action, 2*(1), 2005.
Jha, A. K., Bloch, R., & Lamond, J. (2012). *Cities and flooding: A guide to integrated urban flood risk management for the 21st century*. The World Bank.
Kay, J. E., Deser, C., Phillips, A., Mai, A., Hannay, C., Strand, G., & Holland, M. (2015). The community earth system model (CESM) large ensemble project: A community resource for studying climate change in the presence of internal climate variability. *Bulletin of the American Meteorological Society, 96*(8), 1333–1349.
Kendon, E. J., Roberts, N. M., Fowler, H. J., Roberts, M. J., Chan, S. C., & Senior, C. A. (2014). Heavier summer downpours with climate change revealed by weather forecast resolution model. *Nature Climate Change, 4*(7), 570–576.
Ketterer, C., & Matzarakis, A. (2014). Human-biometeorological assessment of the urban heat island in a city with complex topography–the case of Stuttgart, Germany. *Urban Climate, 10*, 573–584.
Kim, Y., Smith, J. B., Mack, C., Cook, J., Furlow, J., Njinga, J. L., & Cote, M. (2017). A perspective on climate-resilient development and national adaptation planning based on USAID's experience. *Climate and Development, 9*(2), 141–151.
Kinney, P. L., Schwartz, J., Pascal, M., Petkova, E., Le Tertre, A., Medina, S., & Vautard, R. (2015). Winter season mortality: Will climate warming bring benefits? *Environmental Research Letters, 10*(6), 064016.
Kjellstrom, T., Holmer, I., & Lemke, B. (2009). Workplace heat stress, health and productivity â€an increasing challenge for low and middle-income countries during climate change. *Global Health Action, 2*.

ARTICLE IN PRESS

Kovarik, K. R., & Kovarik, J. E. 2016. U.S. Patent No. 9,408,880. Washington, DC: U.S. Patent and Trademark Office.
Kraas, F. (2008). Megacities as global risk areas. *Urban Ecology,* 583–596.
Krantz, L. (2001). *The sustainable livelihood approach to poverty reduction.* SIDA. Division for Policy and Socio-Economic Analysis.
Kruger, A. C., & Sekele, S. S. (2013). Trends in extreme temperature indices in South Africa: 1962–2009. *International Journal of Climatology, 33*(3), 661–676.
Kuenzer, C., & Renaud, F. G. (2012). Climate and environmental change in river deltas globally: Expected impacts, resilience, and adaptation. *The Mekong Delta system* (pp. 7–46). Springer Netherlands.
Kundzewicz, Z. W. (2016). *Extreme weather events and their consequences*.
Landauer, M., Juhola, S., & Klein, J. (2018). The role of scale in integrating climate change adaptation and mitigation in cities. *Journal of Environmental Planning and Management,* 1–25.
Lavda, A. C. (2010). *Psychological factors in skin conditions* (Doctoral dissertation) University of Sheffield.
Lema, M. A., & Majule, A. E. (2009). Impacts of climate change, variability and adaptation strategies on agriculture in semi-arid areas of Tanzania: The case of Manyoni District in Singida Region, Tanzania. *African Journal of Environmental Science and Technology, 3*(8), 206–218.
Levy, K., Woster, A. P., Goldstein, R. S., & Carlton, E. J. (2016). Untangling the impacts of climate change on waterborne diseases: A systematic review of relationships between diarrheal diseases and temperature, rainfall, flooding, and drought. *Environmental Science & Technology, 50*(10), 4905–4922.
Lin, T. P., de Dear, R., & Hwang, R. L. (2011). Effect of thermal adaptation on seasonal outdoor thermal comfort. *International Journal of Climatology, 31*(2), 302–312.
Lo, C. P., & Quattrochi, D. A. (2003). Land-use and land-cover change, urban heat island phenomenon, and health implications. *Photogrammetric Engineering & Remote Sensing, 69*(9), 1053–1063.
Lorenzini, G., Nali, C., & Pellegrini, E. (2014). Summer heat waves, agriculture, forestry and related issues: An introduction. *Agrochimica, 58*(1), 3–19.
Lukamba, M. T. (2010). Natural disasters in African countries: What can we learn about them? *TD: The Journal for Transdisciplinary Research in Southern Africa, 6*(2), 478–495.
Lung, T., Lavalle, C., Hiederer, R., Dosio, A., & Bouwer, L. M. (2013). A multi-hazard regional level impact assessment for Europe combining indicators of climatic and non-climatic change. *Global Environmental Change, 23*(2), 522–536.
Macdonald, R. W., Mackay, D., Li, Y. F., & Hickie, B. (2003). How will global climate change affect risks from long-range transport of persistent organic pollutants? *Human and Ecological Risk Assessment, 9*(3), 643–660.
Mahmood, R., Pielke, R. A., Hubbard, K. G., Niyogi, D., Dirmeyer, P. A., McAlpine, C., ... Baker, B. (2014). Land cover changes and their biogeophysical effects on climate. *International Journal of Climatology, 34*(4), 929–953.
Matysiak, A., & Roess, A. (2017). Interrelationship between climatic, ecologic, social, and cultural determinants affecting dengue emergence and transmission in Puerto Rico and their implications for Zika response. *Journal of Tropical Medicine, 2017*.
McGeehin, M. A., & Mirabelli, M. (2001). The potential impacts of climate variability and change on temperature-related morbidity and mortality in the United States. *Environmental Health Perspectives, 109*(Suppl. 2), 185.
McGranahan, G., Balk, D., & Anderson, B. (2007). The rising tide: Assessing the risks of climate change and human settlements in low elevation coastal zones. *Environment and Urbanization, 19*(1), 17–37.
McMichael, A. J. (2014). Earth as humans' habitat: Global climate change and the health of populations. *International Journal of Health Policy and Management, 2*(1), 9.
McMichael, A. J., Wilkinson, P., Kovats, R. S., Pattenden, S., Hajat, S., Armstrong, B., & Kosnik, M. (2008). International study of temperature, heat and urban mortality: The 'ISOTHURM' project. *International Journal of Epidemiology, 37*(5), 1121–1131.
Mellor, J. E., Levy, K., Zimmerman, J., Elliott, M., Bartram, J., Carlton, E., & Lantagne, D. (2016). Planning for climate change: The need for mechanistic systems-based approaches to study climate change impacts on diarrheal diseases. *Science of the Total Environment, 548*, 82–90.
Metzger, K. B., Ito, K., & Matte, T. D. (2010). Summer heat and mortality in New York City: How hot is too hot? *Environmental Health Perspectives, 118*(1), 80.
Mitchell, B. C., & Chakraborty, J. (2015). Landscapes of thermal inequity: Disproportionate exposure to urban heat in the three largest US cities. *Environmental Research Letters, 10*(11), 115005.
Moore, T. H., Zammit, S., Lingford-Hughes, A., Barnes, T. R., Jones, P. B., Burke, M., & Lewis, G. (2007). Cannabis use and risk of psychotic or affective mental health outcomes: A systematic review. *The Lancet, 370*(9584), 319–328.
Moran, D. S., & Epstein, Y. (2006). Evaluation of the environmental stress index (ESI) for hot/dry and hot/wet climates. *Industrial Health, 44*(3), 399–403.
Moritz, C., & Agudo, R. (2013). The future of species under climate change: Resilience or decline? *Science, 341*(6145), 504–508.
Mutunga, C., & Hardee, K. (2010). Population and reproductive health in National Adaptation Programmes of Action (NAPAs) for climate change in Africa. *African journal of reproductive health, 14*(4).
Nadin, R., Opitz-Stapleton, S., & Yinlong, X. (2015). *Climate risk and resilience in China.* Routledge.
Nehama, F. P. J., Matavel, A. J., Hoguane, A. M., Menomussanga, M., Hoguane, C. A. M., Zacarias, O., & Lemos, M. A. (2016). Building community resilience and strengthening local capacities for disaster risk reduction and climate change adaptation in Zongoene (Xai-Xai District), Gaza Province. *Climate change and health* (pp. 369–385). Springer International Publishing.
Neira, M., & Prüss-Ustün, A. (2016). Preventing disease through healthy environments: A global assessment of the environmental burden of disease. *Toxicology Letters, 259*, S1.
Nguyen, T. T., Bonetti, J., Rogers, K., & Woodroffe, C. D. (2016). Indicator-based assessment of climate-change impacts on coasts: A review of concepts, methodological approaches and vulnerability indices. *Ocean and Coastal Management, 123*, 18–43.
Nkonya, E., Place, F., Kato, E., & Mwanjololo, M. (2015). Climate risk management through sustainable land management in sub-Saharan Africa. *Sustainable intensification to advance food security and enhance climate resilience in Africa* (pp. 75–111). Cham: Springer.
Ogunjobi, K. O., Adamu, Y., Akinsanola, A. A., & Orimoloye, I. R. (2018). Spatiotemporal analysis of land use dynamics and its potential indications on land surface temperature in Sokoto Metropolis, Nigeria. *Royal Society Open Science, 5*, 180661.
Oka, M. (2011). The influence of urban street characteristics on pedestrian heat comfort levels in Philadelphia. *Transactions in GIS, 15*(1), 109–123.
Onamuti, O. Y., Okogbue, E. C., & Orimoloye, I. R. (2017). Remote sensing appraisal of Lake Chad shrinkage connotes severe impacts on green economics and socio-economics of the catchment area. *Royal Society Open Science, 4*(11), 171120.
O'Neill, M. S., & Ebi, K. L. (2009). Temperature extremes and health: Impacts of climate variability and change in the United States. *Journal of Occupational and Environmental Medicine, 51*(1), 13–25.
Onozuka, D., Hashizume, M., & Hagihara, A. (2010). Effects of weather variability on infectious gastroenteritis. *Epidemiology and Infection, 138*(2), 236–243.
Onyango, E. A., Sahin, O., Awiti, A., Chu, C., & Mackey, B. (2016). An integrated risk and vulnerability assessment framework for climate change and malaria transmission in East Africa. *Malaria Journal, 15*(1), 551.
Opiyo, E. O. (2014). *Climate variability and change on vulnerability and adaptation among turukana pastoralist in north-western Kenya* (A PhD thesis submitted to the Department of Rangeland management)University of Nairobi.
Orimoloye, I. R., & Adigun, A. I. (2017). Response of cassava and maize yield to varying spatial scales of rainfall and temperature scenarios in Port Harcourt. *Research Journal of Environmental Sciences, 11*, 137–142.
Orimoloye, I. R., Mazinyo, S. P., Nel, W., & Iortyom, E. T. (2018). *Assessing changes in climate variability observation and simulation of temperature and relative humidity: A case of East London, South Africa*.
Orimoloye, I. R., Mazinyo, S. P., Nel, W., & Kalumba, A. M. (2018). Spatiotemporal monitoring of land surface temperature and estimated radiation using remote sensing: Human health implications for East London, South Africa. *Environmental Earth Sciences, 77*(3), 77.
Otto-Bliesner, B. L., Brady, E. C., Fasullo, J., Jahn, A., Landrum, L., Stevenson, S., ... Strand, G. (2016). Climate variability and change since 850 CE: An ensemble approach with the community earth system model. *Bulletin of the American Meteorological Society, 97*(5), 735–754.
Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature, 421*(6918), 37–42.
Parsons, K. (2014). *Human thermal environments: The effects of hot, moderate, and cold environments on human health, comfort, and performance.* CRC press.
Patz, J. A., & Frumkin, H. (2016). Climate change and human health. *Environmental health: From global to local* (pp. 275). .
Perry, A. G., Korenberg, M. J., Hall, G. G., & Moore, K. M. (2011). Modeling and syndromic surveillance for estimating weather-induced heat-related illness. *Journal of environmental and public health, 2011*.
Phung, D., Guo, Y., Thai, P., Rutherford, S., Wang, X., Nguyen, M., & Chu, C. (2016). The effects of high temperature on cardiovascular admissions in the most populous tropical city in Vietnam. *Environmental Pollution, 208*, 33–39.
Phung, D., Huang, C., Rutherford, S., Chu, C., Wang, X., Nguyen, M., & Nguyen, T. H. (2015). Association between climate factors and diarrhoea in a Mekong Delta area. *International Journal of Biometeorology, 59*(9), 1321–1331.
Portier, C. J., Thigpen, T. K., Carter, S. R., Dilworth, C. H., Grambsch, A. E., Gohlke, J., ... Maslak, T. (2017). *A human health perspective on climate change: A report outlining the research needs on the human health effects of climate change.* Environmental Health Perspectives/National Institute of Environmental Health Sciences.
Quan, J., Chen, Y., Zhan, W., Wang, J., Voogt, J., & Wang, M. (2014). Multi-temporal trajectory of the urban heat island centroid in Beijing, China based on a Gaussian volume model. *Remote Sensing of Environment, 149*, 33–46.
Queiroz, M. P. G., Naas, I. D. A., & Sampaio, C. A. D. P. (2005). *Estimating thermal comfort for piglets considering ammonia concentration*.
Reason, C. J. C., & Keibel, A. (2004). Tropical cyclone Eline and its unusual penetration and impacts over the southern African mainland. *Weather and Forecasting, 19*(5), 789–805.
Reckien, D., Flacke, J., Dawson, R. J., Heidrich, O., Olazabal, M., Foley, A., ... Geneletti, D. (2014). Climate change response in Europe: what's the reality? Analysis of adaptation and mitigation plans from 200 urban areas in 11 countries. *Climatic Change, 122*(1–2), 331–340.
Reed, M. S., Podesta, G., Fazey, I., Geeson, N., Hessel, R., Hubacek, K., & Ritsema, C. (2013). Combining analytical frameworks to assess livelihood vulnerability to climate change and analyse adaptation options. *Ecological Economics, 94*, 66–77.
Rehfuess, E. A., & Bartram, J. (2014). Beyond direct impact: Evidence synthesis towards a better understanding of effectiveness of environmental health interventions. *International Journal of Hygiene and Environmental Health, 217*(2), 155–159.
Rivas, M., Araya, M. C., Caba, F., Rojas, E., & Calaf, G. M. (2011). Ultraviolet light exposure influences skin cancer in association with latitude. *Oncology Reports, 25*(4), 1153–1159.
Ropo, O. I., Perez, M. S., Werner, N., & Enoch, T. I. (2017). Climate variability and heat stress index have increasing potential Ill-health and environmental impacts in the East London, South Africa. *International Journal of Applied Engineering Research, 12*(17), 6910–6918.
Rosenzweig, C., & Iglesias, A. (1994). *Implications of climate change for international agriculture*.
Rosenzweig, C., Solecki, W. D., Romero-Lankao, P., Mehrotra, S., Dhakal, S., & Ibrahim, S. A. (2018). *Climate change and cities: Second assessment report of the urban climate*

change research network. Cambridge University Press.
Rouhani, P., Pinheiro, P. S., Sherman, R., Arheart, K., Fleming, L. E., Mackinnon, J., & Kirsner, R. S. (2010). Increasing rates of melanoma among nonwhites in Florida compared with the United States. *Archives of Dermatology, 146*(7), 741–746.
Rounsevell, M. D. A., Dawson, T. P., & Harrison, P. A. (2010). A conceptual framework to assess the effects of environmental change on ecosystem services. *Biodiversity and Conservation, 19*(10), 2823–2842.
Ryti, N. R., Guo, Y., & Jaakkola, J. J. (2015). Global association of cold spells and adverse health effects: A systematic review and meta-analysis. *Environmental Health Perspectives, 124*(1), 12–22.
Sahu, S., Sett, M., & Kjellstrom, T. (2013). Heat exposure, cardiovascular stress and work productivity in rice harvesters in India: Implications for a climate change future. *Industrial Health, 51*(4), 424–431.
Satterthwaite, D. (2007). *Adapting to climate change in urban areas: the possibilities and constraints in low-and middle-income nations. Vol. 1*. Iied.
Schoen, C. (2005). A new empirical model of the temperature–humidity index. *Journal of applied meteorology, 44*(9), 1413–1420.
Fischer, E. M., & Schär, C. (2010). Consistent geographical patterns of changes in high-impact European heatwaves. *Nature Geoscience, 3*(6), 398–403.
Schipper, E. L. F. (2009). Meeting at the crossroads?: Exploring the linkages between climate change adaptation and disaster risk reduction. *Climate and Development, 1*(1), 16–30.
Schrempf, M., Haluza, D., Simic, S., Riechelmann, S., Graw, K., & Seckmeyer, G. (2016). Is multidirectional UV exposure responsible for increasing melanoma prevalence with altitude? A hypothesis based on calculations with a 3D-human exposure model. *International Journal of Environmental Research and Public Health, 13*(10), 961.
Schwartz, J., Samet, J. M., & Patz, J. A. (2004). Hospital admissions for heart disease: The effects of temperature and humidity. *Epidemiology, 15*(6), 755–761.
Seaman, J. A., Sawdon, G. E., Acidri, J., & Petty, C. (2014). The household economy approach. Managing the impact of climate change on poverty and food security in developing countries. *Climate Risk Management, 4*, 59–68.
Seckmeyer, G., Schrempf, M., Wieczorek, A., Riechelmann, S., Graw, K., Seckmeyer, S., & Zankl, M. (2013). A novel method to calculate solar UV exposure relevant to vitamin D production in humans. *Photochemistry and Photobiology, 89*(4), 974–983.
Senapati, S., & Gupta, V. (2017). Socio-economic vulnerability due to climate change: Deriving indicators for fishing communities in Mumbai. *Marine Policy, 76*, 90–97.
Sewankambo, N., Tumwine, J. K., & Besada, H. (2016). The impact of climate change on health governance and sovereignty in Africa. *Moving health sovereignty in Africa: Disease, governance, climate change* (pp. 167). .
Seyfried, T. N. (2015). Cancer as a mitochondrial metabolic disease. *Frontiers in Cell and Development Biology, 3*.
Shamsuddoha, M., Ullah, M. S., & Shahjahan, M. (2014). 5 land acquisition for climate-displaced communities in Bangladesh. *Land Solutions for Climate Displacement,* 159.
Shanklin, J. (2004). Book review: Global warming: The complete briefing, (Houghton). *The Journal of the British Astronomical Association, 114*, 362.
Shaw, A. (2018). Towards sustainable cities in India. *Sustainable urbanization in India* (pp. 23–37). Singapore: Springer.
Shi, L., Kloog, I., Zanobetti, A., Liu, P., & Schwartz, J. D. (2015). Impacts of temperature and its variability on mortality in New England. *Nature Climate Change, 5*(11), 988–991.
Singh, R. B., Hales, S., De Wet, N., Raj, R., Hearnden, M., & Weinstein, P. (2001). The influence of climate variation and change on diarrheal disease in the Pacific Islands. *Environmental Health Perspectives, 109*(2), 155.
Smoyer-Tomic, K. E., & Rainham, D. G. (2001). Beating the heat: Development and evaluation of a Canadian hot weather health-response plan. *Environmental Health Perspectives, 109*(12), 1241.
Sonwa, D. J., Somorin, O. A., Jum, C., Bele, M. Y., & Nkem, J. N. (2012). Vulnerability, forest- related sectors and climate change adaptation: The case of Cameroon. *Forest Policy and Economics, 23*, 1–9.
Soret, S., Mejia, A., Batech, M., Jaceldo-Siegl, K., Harwatt, H., & Sabaté, J. (2014). Climate change mitigation and health effects of varied dietary patterns in real-life settings throughout North America. *The American Journal of Clinical Nutrition, 100*(Supplement 1), 490S–495S.
Steadman, R. G. (1979). The assessment of sultriness. Part I: A temperature-humidity index based on human physiology and clothing science. *Journal of Applied Meteorology, 18*(7), 861–873.
Stephen, H., Iortyom, E. T., Ropo, O. I., & Daniel, D. P. (2017). Analysis of the physical growth and expansion of Makurdi town using remote sensing and GIS techniques. *Imperial Journal of Interdisciplinary Research, 3*(7).
Streutker, D. R. (2003). Satellite-measured growth of the urban heat island of Houston, Texas. *Remote Sensing of Environment, 85*(3), 282–289.
Stringer, L. C., Dyer, J. C., Reed, M. S., Dougill, A. J., Twyman, C., & Mkwambisi, D. (2009). Adaptations to climate change, drought and desertification: Local insights to enhance policy in southern Africa. *Environmental Science & Policy, 12*(7), 748–765.
Stull, R. (2000). *Meteorology for Scientists and Engineers* (Second Edition). Brooks/Cole978053437214960.
Suomi, J. (2014). *Characteristics of urban heat island (UHI) in a high latitude coastal city-a case study of Turku, SW Finland*.
Tan, J., Zheng, Y., Tang, X., Guo, C., Li, L., Song, G., & Chen, H. (2010). The urban heat island and its impact on heat waves and human health in Shanghai. *International Journal of Biometeorology, 54*(1), 75–84.
Tietenberg, T. H., & Lewis, L. (2016). *Environmental and natural resource economics*. Routledge.
Tirado, M. C., Cohen, M. J., Aberman, N., Meerman, J., & Thompson, B. (2010). Addressing the challenges of climate change and biofuel production for food and nutrition security. *Food Research International, 43*(7), 1729–1744.
Trnka, M., Olesen, J. E., Kersebaum, K. C., Skjelvåg, A. O., Eitzinger, J., Seguin, B., ... Dubrovský, M. (2011). Agroclimatic conditions in Europe under climate change. *Global Change Biology, 17*(7), 2298–2318.
Turner, W. R., Bradley, B. A., Estes, L. D., Hole, D. G., Oppenheimer, M., & Wilcove, D. S. (2010). Climate change: Helping nature survive the human response. *Conservation Letters, 3*(5), 304–312.
Tursilowati, L., & Djundjunan, J. D. (2007). Use of remote sensing and GIS to compute temperature humidity index as human comfort indicator relate with land use-land cover change (LULC) in Surabaya. *Proceedings of the 73rd international symposium on sustainable humanosphere, Bandung, Indonesia* (pp. 22–25). .
Vaneckova, P., Hart, M. A., Beggs, P. J., & De Dear, R. J. (2008). Synoptic analysis of heat-related mortality in Sydney, Australia, 1993–2001. *International Journal of Biometeorology, 52*(6), 439–451.
Vaneckova, P., Neville, G., Tippett, V., Aitken, P., FitzGerald, G., & Tong, S. (2011). Do biometeorological indices improve modeling outcomes of heat-related mortality? *Journal of Applied Meteorology and Climatology, 50*(6), 1165–1176.
Vargo, J., Stone, B., Habeeb, D., Liu, P., & Russell, A. (2016). The social and spatial distribution of temperature-related health impacts from urban heat island reduction policies. *Environmental Science & Policy, 66*, 366–374.
Venton, P., & La Trobe, S. (2008). Linking climate change adaptation and disaster risk reduction. *Linking climate change adaptation and disaster risk reduction*. Tearfund; Institute of Development Studies (IDS).
Von-Doellinger, V., Campos, M., Mendes, L., & Schramm, J. (2014). The 2008 Global Burden of Disease study in Brazil: a new methodological approach for estimation of injury morbidity. *Revista Panamericana de Salud Pública, 36*(6), 368–375.
Voogt, J. A., & Oke, T. R. (2003). Thermal remote sensing of urban climates. *Remote Sensing of Environment, 86*(3), 370–384.
Voskamp, I. M., & Van de Ven, F. H. M. (2015). Planning support system for climate adaptation: Composing effective sets of blue-green measures to reduce urban vulnerability to extreme weather events. *Building and Environment, 83*, 159–167.
Waas, T., & Hugé, J. (2012). Developing an environmental sustainability toolkit to integrate climate change issues in development cooperation. *Climate change and the sustainable use of water resources* (pp. 401–413). Springer Berlin Heidelberg.
Walther, G. R. (2010). Community and ecosystem responses to recent climate change. *Philosophical Transactions of the Royal Society of London B, 365*(1549), 2019–2024.
Wang, T., Hamann, A., Spittlehouse, D. L., & Murdock, T. Q. (2012). ClimateWNA—high-resolution spatial climate data for western North America. *Journal of Applied Meteorology and Climatology, 51*(1), 16–29.
Watson, J. E. (2014). Human responses to climate change will seriously impact biodiversity conservation: it's time we start planning for them. *Conservation Letters, 7*(1), 1–2.
Weihs, P., Schmalwieser, A., Reinisch, C., Meraner, E., Walisch, S., & Harald, M. (2013). Measurements of personal UV exposure on different parts of the body during various activities. *Photochemistry and Photobiology, 89*(4), 1004–1007.
Weisskopf, M. G., Anderson, H. A., Foldy, S., Hanrahan, L. P., Blair, K., Török, T. J., & Rumm, P. D. (2002). Heat wave morbidity and mortality, Milwaukee, Wis, 1999 vs 1995: An improved response? *American Journal of Public Health, 92*(5), 830–833.
Wright, C. Y., Street, R. A., Cele, N., Kunene, Z., Balakrishna, Y., Albers, P. N., & Mathee, A. (2017). Indoor temperatures in patient waiting rooms in eight rural primary health care centers in northern South Africa and the related potential risks to human health and wellbeing. *International Journal of Environmental Research and Public Health, 14*(1), 43.
Xiang, J., Bi, P., Pisaniello, D., & Hansen, A. (2014). Health impacts of workplace heat exposure: An epidemiological review. *Industrial Health, 52*(2), 91–101.
Xu, H., Hu, X., Guan, H., & He, G. (2017). Development of a fine-scale discomfort index map and its application in measuring living environments using remotely-sensed thermal infrared imagery. *Energy and Buildings, 150*, 598–607.
Yawson, D. O., Adu, M. O., Armah, F. A., Kusi, J., Ansah, I. G., & Chiroro, C. (2015). A needs-based approach for exploring vulnerability and response to disaster risk in rural communities in low income countries. *Australasian Journal of Disaster and Trauma Studies, 19*, 27.
Yu, W., Vaneckova, P., Mengersen, K., Pan, X., & Tong, S. (2010). Is the association between temperature and mortality modified by age, gender and socio-economic status? *Science of the Total Environment, 408*(17), 3513–3518.
Zanobetti, A., & Schwartz, J. (2005). The effect of particulate air pollution on emergency admissions for myocardial infarction: a multicity case-crossover analysis. *Environmental health perspectives, 113*(8), 978.
Ziervogel, G., New, M., Archer van Garderen, E., Midgley, G., Taylor, A., Hamann, R., & Warburton, M. (2014). Climate change impacts and adaptation in South Africa. *Wiley Interdisciplinary Reviews: Climate Change, 5*(5), 605–620.